

Strange Attractor in the Potts Spin Glass on Hierarchical Lattices [☆]

Washington de Lima^a, G. Camelo-Neto^b, S. Coutinho^{c,*}

^aUniversidade Federal de Pernambuco, Campus Caruaru, Pernambuco, Brazil.

^bUniversidade Federal de Alagoas, Núcleo de Ciências Exatas, Laboratório de Física Teórica e Computacional, CEP 57309-005, Arapiraca, Alagoas, Brazil.

^cUniversidade Federal de Pernambuco, Departamento de Física, Laboratório de Física Teórica e Computacional, Cidade Universitária, CEP 50670-901, Recife, Pernambuco, Brazil.

Abstract

The spin-glass q -state Potts model on d -dimensional diamond hierarchical lattices is investigated by an exact real space renormalization group scheme. Above a critical dimension $d_l(q)$ for $q > 2$, the coupling constants probability distribution flows to a low temperature *strange attractor* or to the high temperature paramagnetic fixed point, according to the temperature is below or above the critical temperature $T_c(q, d)$. The strange attractor was investigated considering four initial different distributions for $q = 3$ and $d = 5$ presenting strong robustness in shape and temperature interval suggesting a condensed phase with algebraic decay.

Keywords: Spin Glass, Potts Model, Strange Attractor, Hierarchical Lattices.

PACS: 05.50. +q, 64.60. Ak, 75.10. Nr

1. Introduction

The q -states Potts model, proposed a long time ago by Domb as the subject of Potts doctoral thesis [1], has found a wide range of applicability in many fields of both basic and material sciences. The Potts model was conceived as a generalization of the Ising model [2], when $q = 2$, and the Askin-Teller model ($q = 4$) [3]. It also mimics the problem of percolation ($q = 1$) [4, 5] and even the problem of the linear resistor networks ($q = 0$) [6]. All of the above mentioned problems were also encompassed by the random-cluster model introduced by Fortuin and Kasteleyn [7]. Furthermore, the degeneracy of the ground state of the *antiferromagnetic* Potts models was shown to be related with the q -coloring problem [7]. It is important to emphasize that the most important feature of the mathematical structure of the Potts model is the equivalence between its partition function and Tutte polynomial [8]. Concerning applications, the Potts model has been applied in many fields, such as biology [9], sociology [10] and material science [11]. In the latter, for instance, the technique of Monte Carlo simulations on the Potts model has been applied to a wide variety of phenomena, such as diffusion in polycrystalline microstructures [12] and the study of viscous instabilities in foam-flow behavior [13].

In this letter, the properties of the q -state Potts model with random competing interactions are investigated. This

model is called Potts glass in allusion to the particular case when $q = 2$, widely known in the literature as the Ising spin-glass model. The absence of spin-inversion symmetry and a different nature of the frustration effects distinguish it from its Ising counterpart, exhibiting rather a richer critical behavior in mean-field theory [14, 15, 16]. The pure and disordered Potts model (without frustration) and related models have been widely investigated in the past. For review papers see, for instance, [6, 17]. The present work focuses on the study of the q -state Potts model with random frustrated exchange interactions on a family of diamond-type hierarchical lattices [18] with scale factor $b = 2$. When symmetrical zero-centered random exchange couplings distributions are considered the system undergoes a phase transition from a paramagnetic high temperature phase to a low-temperature condensed phase above some dimension, $d_l(q)$. The phase diagram of the Potts glass model had been investigated before, by the Migdal-Kadanoff (M-K) renormalization group (RG) scheme, indicating the presence of a condensed phase at finite temperatures when $q > 2$ and $d = 4$ [19]. In reference [19], however, it was assumed (as working *hypothesis*) that the initial symmetrical Gaussian probability distribution of coupling constants with variance σ is transformed under renormalization into another symmetrical Gaussian distribution but with variance $\sqrt{b^{d-1}}\sigma$.

The M-K real-space RG method for Bravais lattices is known to be equivalent to exactly solving the model on diamond-like hierarchical lattices [20, 21]. In a recent paper [22], however, the authors studied the occurrence of phase transitions of the Potts glass model using this exact approach by numerically following the flow of the

[☆]Work supported by the Brazilian granting agencies CNPq (grant PRONEX/FACEPE 0203-1.05/08), and CAPES

*Corresponding author: S. Coutinho

Email address: sergio@lftc.ufpe.br (S. Coutinho)

renormalized probability distribution in an appropriated parameter space. Such space was previously considered in reference [23, 24] to study the Ising spin-glass model. The q -state Potts glass model was considered on lattices with several fractal dimension, determining the critical temperature and the upper and lower bounds for the associated lower critical dimension $d_l(q)$. For instance, for $q = 3$ the lower (upper) bound was found to be 4.46 (4.58), in contrast with the result obtained in reference[19], which finds the transition occurring for $d < 4$. Here we further explore the flow of the renormalized probability distribution in the whole parameter space and investigate the nature of the low temperature stable fixed point, which surprisingly appeared like a *strange attractor*.

2. Renormalization Procedure in Disordered Systems

For pure systems, the renormalization procedure consists in finding the equivalent exchange interaction for a pair of spins after eliminating several spins in the lattice. For a disordered system, however, the renormalization procedure will affect the whole distribution of coupling constants, the renormalized distribution, $P'(J)$, is related with the previous (non-renormalized) distribution, $P(J)$, by,

$$P'(J) = \int \cdots \int \prod_{\langle ij \rangle} P(J_{ij}) dJ_{ij} \delta(K - K'(K)), \quad (1)$$

where $K = \beta J$ is a reduced coupling constant, $K'(K)$ is the renormalization equation, and the product runs over all the pairs of spins $\langle ij \rangle$.

Equation (1) should be iterated until the renormalized distribution reaches a fixed point distribution, characteristic of the thermodynamic phase. A zero-centered Dirac-delta distribution, for instance, indicates a paramagnetic phase. The procedure adopted in this work is to produce a sample of random coupling constants from an initial probability density function, feed the renormalization equation to find a sample of the same size of renormalized coupling constants, estimating the physical quantities numerically from the samples. The process is repeated until the fixed point distribution is reached.

3. The Potts Hamiltonian and the Renormalization Equation

The Potts Hamiltonian is written as,

$$\mathcal{H} = - \sum_{\langle ij \rangle} q J_{ij} \delta_{\sigma_i \sigma_j}, \quad (2)$$

where the sum is taken over all bonds in the lattice, the Kronecker δ symbol takes the values 1 if $\sigma_i = \sigma_j$ or 0 otherwise, and $\sigma_i = 1, 2, \dots, q$ are the q -states Potts spins variables, located at each site of a diamond-type hierarchical lattice with p branches and scale factor b . The basic




Figure 1: Renormalization group scheme on the d_f dimension diamond-type hierarchical lattice with p branches and scaling factor $b = 2$.

unit of such lattice is illustrated in figure 1, where μ and μ' are called external sites and the set $\{\sigma_i\}$ represents the internal sites [18]. K_i and L_i are reduced coupling constants, $K_i \equiv \beta J_{\mu\sigma_i}$ and $L_i \equiv \beta J_{\mu'\sigma_i}$. The lattice generations or hierarchies are successively built by replacing each connection of the basic unit by the basic unit itself, yielding to a graph with fractal dimension,

$$d_f = 1 + \frac{\ln p}{\ln 2}.$$

The exact renormalization process on a n -generation lattice consists in partially tracing the partition function along all the internal sites introduced in the n^{th} generation leading to a $(n-1)$ -generation lattice with a set of effective reduced coupling constants $\{K'_i\}$ given by

$$K' = \frac{1}{q} \sum_{i=1}^p \left[\frac{(q-1) + \exp(q K_i + q L_i)}{(q-2) + \exp(q K_i) + \exp(q L_i)} \right]. \quad (3)$$

Equation (3) is the local renormalization equation. The right-hand side of equation (3) receives values for the reduced coupling constants calculated from the n -generation of the coupling constant distribution, $P(J)$, resulting in one of the possible values of the renormalized reduced coupling constant of the $(n-1)$ -generation. The renormalization procedure starts from the thermodynamic limit (generation $n \rightarrow \infty$) where the coupling constants are assumed to have a well known distribution, actually, Gaussian, Delta-Bimodal, Uniform, or Exponential, and well defined temperature T .

3.1. Probability distribution renormalization flow

For each n -generation lattice we can define a set of *thermal transmissivities* variables $\{t_{ij}\}$, each one associated with the respective bond, ie.,

$$t_{ij} \equiv \frac{1 - \exp(-q\beta J_{ij})}{1 + (q-1) \exp(-q\beta J_{ij})}. \quad (4)$$

Thermal transmissivity t_{ij} represents the pair correlation function Γ_{ij} between sites ij [17].

A system with a probability distribution of coupling constants $P(J) \equiv P(\{J_{ij}\})$ yields a thermal transmissivity variance,

$$\Delta^2 \equiv [(t_{ij} - [t_{ij}]_J)^2]_J, \quad (5)$$

where $[\dots]_J$ means the average over the probability distribution, $P(J)$.

Every fixed probability distribution has a signature in the diagram $\Delta^2 \times T$ when the temperature is varied. In the case of the initial probability distributions considered in this work such representation can be seen in figure 2 for the case $q = 3$. Notice that the $T \rightarrow 0$ limit is the




Figure 2: Δ^2 behavior for distinct probabilities distributions against temperature for the $q = 3$ case.

same for all cases indicating an universal zero temperature behavior. Such universal point $(\Delta^{*2}, T = 0)$ can be analytically obtained in the case of the Bimodal distribution, $P(J_{ij}) = [\delta(J_{ij} - 1) + \delta(J_{ij} + 1)]/2$, ie.

$$\Delta^{*2} = \left[\frac{q}{2(q-1)} \right]^2. \quad (6)$$

Note that for $q = 2$, the Ising limit case, $\Delta^* \rightarrow 1$ is recovered as seen in [23].

For each application of the renormalization equation, an effective temperature proportional to the inverse of the square root of the reduced coupling variance may be defined, ie.

$$T_r \propto \frac{1}{\sqrt{[(K_{ij} - [K_{ij}]_J)^2]_J}}. \quad (7)$$

In this work, the flow of the renormalized probability distributions is followed numerically in the parameter space $\Delta^2 \times T_r$, each point representing a stage of the renormalization process.

3.2. Strange attractor

Four distinct initial symmetrical probability distributions were considered, namely the Gaussian, the delta-bimodal, the uniform and the exponential ones. For lattices with dimension greater than $d_l(q)$ and independently of the nature of the initial distribution, the renormalization flow displays the following same features: the paramagnetic phase is characterized by a *fixed point* at infinity renormalized temperature and zero transmissivity variance, while the low-temperature phase is characterized by a *strange attractor* in the renormalization flux for $q \geq 3$ – or a fixed-point $\Delta = 1, T_r = 0$ for $q = 2$ as previously reported[23] for the Ising SG. The observation of such *strange attractor*, which is located in a region of low but finite temperatures, is the main result to be reported in the present work.

A first signature of the existence of this strange attractor was reported by Banavar and Bray[25] long time ago. These authors observed an unusual behavior in the renormalization of $[K_{ij}]_J$ and its standard deviation in the case $d = 5$ and scale factor $b = 2$ (present case). In their words these quantities “wanders chaotically in a smallish region around $[K_{ij}]_J \sim 14 \dots$ (persisting) for 1000’s of iterations and happens for a wide range of starting temperatures”. Furthermore, they do not give a clear explanation of this behavior but speculate that the delicate balance between energy and entropy causes that wandering phase in $d = 5$.

This study uses the same methodology used in references [23, 25], ie. the successive construction of large pools of renormalized constant couplings, calculated using equation (3) starting from an initial probability distribution and fixed temperature. Figure 3 illustrates the renormal-




Figure 3: Renormalization flow in the $\Delta^2 \times T$ parameter space for the $q = 3$ -Potts model in $d = 5$ dimension for an initial symmetric Gaussian distribution. Dotted curve indicates the low-temperature flow while continuous one illustrates the high-temperature flow, according to the arrows. Inset magnifies the flow bifurcation region.

ization flow diagram in the $\Delta^2 \times T$ space for the $q = 3$ case in $d = 5$ dimensions, starting from a Gaussian distribution of the exchange couplings in two temperatures. For initial temperatures below ~ 3.0 the renormalized distribution of interactions (solid triangles) flows to the strange

attractor located in a finite region of the parameter space, for high initial temperatures ($T \sim 3.5$), the renormalized distribution of interactions (solid circles) flows to high temperatures and zero thermal transmissivity variance, characterizing the paramagnetic phase. Figure 4 displays the strange attractor constructed with 10^5 steps of renormalization, lying in the interval $(0.15, 0.35)$ starting from two distinct initial distributions, (a) the Delta-Bimodal and (b) the Gaussian ones. In reference[22], some of the present




Figure 4: Attractor for two distinct initial distribution, (a) Bimodal and (b) Gaussian

authors, using the same methodology, found the critical temperatures of the model for several lattices, indicating the ones where the transition may occurs for appropriated values of d and q .

The fractal dimension of the attractor was estimated by box counting, yielding to,

$$D = 1.66(1),$$

for every initial distribution and for the case $q = 3$ and $d = 5$. A proper analysis of the attractor is taking place at the moment, by estimating its Lyapunov exponents by the Wolf method [26] with various values of q and d .

The strange attractor describing the condensed phase occurs at a non-zero temperature interval probably due to the existence of residual entropy, such as the unusual

algebraic phase previously studied [27, 28, 29] for the anti-ferromagnetic Potts model. Detailed studies of the correlation functions and the local magnetization are under development in order to determine the nature of such condensed phase. Preliminary studies of the present model on lattices with scale factor $b = 3$, which corresponds to bipartite lattices, indicates however the absence of the strange attractor in any dimension. Moreover, the condensed phase which occurs above a certain lower critical dimension $d_l(q)$ does not presents characteristics of spin-glass phase[30].

4. Conclusions

The q -state spin-glass Potts model undergoes a phase transition whenever $d \geq d_l(q)$, from a high temperature paramagnetic phase to a low temperature condensed phase, characterized by strange attractors. These strange attractors are superimposed in the same region of parameter space independently of the initial distribution of coupling constants and initial temperatures below T_c , even if this initial temperature is chosen close to zero, below the attractor region. Such strange attractors, however, change in shape and position for distinct (q, d) models in lattices with scale factor $b = 2$. The occurrence of the strange attractor in a finite non-zero temperature interval suggests a condensed phase with algebraic decay of the correlation function.

5. Acknowledgment

We gratefuly acknowledge the enlightening and fruitful discussions and comments of E. M. F. Curado, W. A. M. Morgado, A. S. Rosas, and F. D. Nobre. This work received financial support from CNPq and CAPES (Brazilian federal grant agencies), from FACEPE (Pernambuco state grant agency under the grant PRONEX APQ 0203-1.05/08).

References

- [1] R. B. Potts, Some generalized order-disorder transformations, *Mathematical Proceedings of the Cambridge Philosophical Society* 48 (1952) 106–109.
- [2] R. J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press, 1982.
- [3] J. Ashkin, E. Teller, *Statistics of Two-Dimensional Lattices with Four Components*, *Physical Review* 64 (1943) 178–184.
- [4] S. R. Broadbent, J. M. Hammersley, *Percolation processes*, *Mathematical Proceedings of the Cambridge Philosophical Society* 53 (1957) 629.
- [5] P. W. Kasteleyn, C. M. Fortuin, *Journal of the Physical Society of Japan* 26 (Suppl.) (1969) 11.
- [6] F. Wu, *The Potts model*, *Reviews of Modern Physics* 54 (1982) 235.
- [7] C. Fortuin, W. P. Kasteleyn, *On the random-cluster model I. Introduction and relation to other models*, *Physica* 57 (1972) 536–564.
- [8] W. Tutte, *On dichromatic polynomials*, *Journal of Combinatorial Theory* 2 (1967) 301–320.

- [9] L. Sun, Y. F. Chang, X. Cai, A discrete simulation of tumor growth concerning nutrient influence, *International Journal of Modern Physics B* 18 (2004) 2651–2657.
- [10] C. Schulze, Potts-like model for ghetto formation in multicultural societies, *International Journal of Modern Physics C* 16 (2005) 351–355.
- [11] K. G. F. Janssens, D. Raabe, E. Kozeschnik, M. A. Miodownik, B. Nestler, *Computational Materials Engineering: An introduction to microstructure evolution*, Elsevier Academic Press, 2007.
- [12] T. Swiler, V. Tikare, E. Holm, Heterogeneous diffusion effects in polycrystalline microstructures, *Materials Science and Engineering A* 238 (1997) 85–93.
- [13] S. Sanyal, J. A. Glazier, Viscous instabilities in flowing foams: a Cellular Potts Model approach., *Journal of statistical mechanics* (Online) 10008 (2006) p10008.
- [14] D. J. Gross, I. Kanter, H. Sompolinsky, Mean-field theory of the Potts glass, *Physical Review Letters* 55 (1985) 304–307.
- [15] G. Cwiliich, T. R. Kirkpatrick, Mean-field theory and fluctuations in Potts spin glasses. I, *Journal of Physics A: Mathematical and General* 22 (1989) 4971–4987.
- [16] G. Cwiliich, Mean-field theory and fluctuations in Potts spin glasses. II, *Journal of Physics A: Mathematical and General* 23 (1990) 5029–5044.
- [17] C. Tsallis, A. C. N. de Magalhães, Pure and random Potts-like models: Real-space renormalization-group approach, *Physics Reports* 268 (1996) 305.
- [18] R. B. Griffiths, M. Kaufman, Spin systems on hierarchical lattices. Introduction and thermodynamic limit, *Physical Review B* 26 (1982) 5022.
- [19] A. Benyoussef, M. Loulidi, Short-range Potts spin-glass model: Renormalization-group method., *Physical review. B, Condensed matter* 53 (1996) 8215–8218.
- [20] L. P. Kadanoff, Notes on Migdal’s Recursion Formulas, *Annals of Physics* 100 (1976) 359.
- [21] A. A. Migdal, Phase transitions in gauge and spin-lattice systems, *Sov. Phys.-JETP* 42 (1976) 743.
- [22] G. Camelo-Neto, W. De-Lima, S. Coutinho, W. Morgado, Potts spin glass: a renormalization group approach, *Physica A* 342 (2004) 83–89.
- [23] E. M. F. Curado, J. L. Meunier, Spin-Glass in low dimensions and the Migdal-Kadanoff approximation, *Physica A* 149 (1988) 164.
- [24] E. Nogueira Jr., S. Coutinho, F. D. Nobre, E. M. F. Curado, Universality in short-range Ising spin glasses, *Physica A: Statistical Mechanics and its Applications* 271 (1999) 125–132.
- [25] J. R. Banavar, A. J. Bray, Heisenberg and Potts spin glasses: A renormalization-group study, *Physical Review B* 38 (1988) 2564.
- [26] A. Wolf, J. B. Swift, H. L. Swinney, J. A. Vastano, Determining lyapunov exponents from a time series, *Physica D: Nonlinear Phenomena* 16 (1985) 285 – 317.
- [27] A. N. Berker, L. P. Kadanoff, Ground-state entropy and algebraic order at low temperatures, *Journal of Physics A: Mathematical and General* 13 (1980) L259–L264.
- [28] Y. Qin, Z. R. Yang, Diamond-type hierarchical lattices for the Potts antiferromagnet, *Physical Review B* 43 (1991) 8576–8582.
- [29] J. Redinz, A. de Magalhães, E. Curado, Potts antiferromagnetic model on a family of fractal lattices: Exact results for an unusual phase, *Physical Review B* 49 (1994) 6689–6697.
- [30] P. H. A. de Andrade, *Modelo de Potts com Interações Aleatórias em Redes Fractais*, Master’s thesis, Universidade Federal de Pernambuco, 2012.